

Multiple choice questions

- 1) (C), $x^3 - 3^3 = (x-3)(x^2 + 3x + 9)$
- 2) (A), $x = \frac{3 \times 8 + 2 \times -2}{3 + 2} = \frac{20}{5} = 4$
- 3) (B), $x^3 - \sum \alpha \cdot x^2 + \sum \alpha \beta \cdot x - \prod \alpha$
- 4) (C), since D: $-2 \leq x \leq 2$, R: $-\frac{3\pi}{2} \leq y \leq \frac{3\pi}{2}$.
- 5) (D), $6! \times 2$.
- 6) (A), $n = 4, \therefore T = \frac{2\pi}{n} = \frac{\pi}{2}, A^2 = 9, \therefore A = 3$.
- 7) (C), $\int \sin^2 3x dx = \frac{1}{2} \int (1 - \cos 6x) dx$.
- 8) (D), $P(x) = (x+1)(x-3)Q(x) + 2x + 7, P(3) = 13$
- 9) (D), $\frac{d}{dx}(\cos^{-1} ax) = \frac{-a}{\sqrt{1-(ax)^2}}$
- 10) (B), $\angle ABT = 90^\circ - \frac{\theta}{2}$ (angle sum in isos. Δ)
and $\angle APB = \angle ABT$ (alternate segment angles)

Question 11

- (a) $\int_0^3 \frac{1}{9+x^2} dx = \left[\frac{1}{3} \tan^{-1} \frac{x}{3} \right]_0^3 = \frac{1}{3} \times \frac{\pi}{4} = \frac{\pi}{12}$.
- (b) $\frac{d}{dx}(x^2 \tan x) = 2x \tan x + x^2 \sec^2 x$.
- (c) $\frac{x}{x-3} < 2$
 $x(x-3) < 2(x-3)^2$
 $(x-3)(x-2x+6) < 0$
 $(x-3)(-x+6) < 0$
 $x < 3$ or $x > 6$.
- (d) $u = 2 - x, du = -dx$.

When $x = 1, u = 1$; when $x = 2, u = 0$.

$$I = \int_1^0 (2-u)u^5(-du) = \int_0^1 (2u^5 - u^6) du$$

$$= \left[\frac{u^6}{3} - \frac{u^7}{7} \right]_0^1 = \frac{1}{3} - \frac{1}{7} = \frac{4}{21}$$

(e) ${}^8C_3 \times {}^{10}C_4 = 11760$.

(f) (i) $\left(2x^3 - \frac{1}{x} \right)^{12} = \frac{(2x^4 - 1)^{12}}{x^{12}}$.

The constant term is $\frac{{}^{12}C_3 (2x^4)^3 (-1)^9}{x^{12}}$
 $= -{}^{12}C_3 (2)^3 = -1760$.

(ii) $\left(2x^3 - \frac{1}{x} \right)^n = \frac{(2x^4 - 1)^n}{x^n}$.

$$T_{k+1} = \frac{{}^n C_k (2x^4)^k (-1)^{n-k}}{x^n}$$

$\therefore n = 4k, k$ is a non-negative integer.

Question 12

(a) Let $n = 1, 2^3 - 3 = 8 - 3 = 5$, hence, divisible by 5.

Assume $2^{3n} - 3^n = 5M$, where M is an integer.

$$\therefore 2^{3n} = 5M + 3^n.$$

Required to prove that $2^{3(n+1)} - 3^{n+1}$ is divisible by 5.

$$\begin{aligned} 2^{3(n+1)} - 3^{n+1} &= 2^3 \times 2^{3n} - 3 \times 3^n \\ &= 8(5M + 3^n) - 3 \times 3^n \\ &= 40M + 5 \times 3^n \\ &= 5(8M + 3^n), \text{ which is divisible by 5.} \end{aligned}$$

\therefore By the principle of Induction, $2^{3n} - 3^n$ is divisible by 5 for all $n \geq 1$.

(b) (i) $4x - 3 \geq 0, \therefore x \geq \frac{3}{4}$.

(ii) $f : y = \sqrt{4x - 3}$.

$$f^{-1} : x = \sqrt{4y - 3}.$$

$$x^2 = 4y - 3.$$

$$\therefore y = \frac{x^2 + 3}{4}, x \geq 0.$$

(iii) $\frac{x^2 + 3}{4} = x$


$$x^2 - 4x + 3 = 0$$

$$(x - 1)(x - 3) = 0$$

$$\therefore x = 1, 3$$

\therefore Points of intersection (1,1) and (3,3).

(iv)



(c) (i) $\Pr(W) = \Pr(L), \Pr(D) = \frac{1}{5}$.

$$\Pr(W) + \Pr(L) + \Pr(D) = 1.$$

$$\therefore \Pr(W) = \frac{2}{5}.$$

(ii) ${}^6C_3 \left(\frac{2}{5}\right)^3 \left(\frac{3}{5}\right)^3 = 0.2765$.

(d) (i) $AC \perp BC, \therefore \frac{y}{-t} \times \frac{k}{t} = -1, \therefore y = \frac{t^2}{k}$.

$$x_p = x_c = t, y_p = y_b = y = \frac{t^2}{k}.$$

(ii) The equation of the parabola is $y = \frac{x^2}{k}$, i.e.

$$ky = x^2.$$

$$4a = k, \therefore a = \frac{k}{4}.$$

$$\therefore \text{Focus} \left(0, \frac{k}{4}\right).$$

Question 13


$$(a) \sin\left(2\cos^{-1}\frac{2}{3}\right) = \sin 2\alpha = 2\sin\alpha\cos\alpha$$

$$= 2 \times \frac{\sqrt{5}}{3} \times \frac{2}{3} = \frac{4\sqrt{5}}{9}.$$

$$(b) (i) y = \frac{2x^2}{x^2+9} = \frac{2}{1+\frac{9}{x^2}} \rightarrow 2 \text{ as } x \rightarrow \infty.$$

\therefore Horizontal asymptote $y = 2$.

(ii)



$$(c) (i) x = 5 + 6\cos 2t + 8\sin 2t$$

$$= 5 + 10\cos\left(2t - \tan^{-1}\frac{4}{3}\right).$$

$$\dot{x} = -20\sin\left(2t - \tan^{-1}\frac{4}{3}\right).$$

$$\ddot{x} = -40\cos\left(2t - \tan^{-1}\frac{4}{3}\right) = -4(x-5), \therefore \text{SHM.}$$

$$(ii) 0 = 5 + 10\cos\left(2t - \tan^{-1}\frac{4}{3}\right).$$

$$\cos\left(2t - \tan^{-1}\frac{4}{3}\right) = -\frac{1}{2}.$$

$$2t - \tan^{-1}\frac{4}{3} = \frac{2\pi}{3}.$$

$$\therefore t = \frac{1}{2}\tan^{-1}\frac{4}{3} + \frac{\pi}{3} \text{ sec} \approx 1.5 \text{ sec.}$$

$$(d) (i) \frac{dC}{dt} = 1.4(e^{-0.2t} - 0.2te^{-0.2t}) = 1.4e^{-0.2t}(1 - 0.2t)$$

$$\frac{dC}{dt} = 0 \text{ gives } t = \frac{1}{0.2} = 5 \text{ hours.}$$

$$\frac{d^2C}{dt^2} = 1.4(-0.2e^{-0.2t}(1 - 0.2t) - 0.2e^{-0.2t})$$

$$= -0.28e^{-0.2t}(2 - 0.2t) < 0 \text{ when } t = 5.$$

\therefore The maximum occurs when $t = 5$ hours.

$$(ii) C_2 = 20 + \frac{0.3 - 1.4 \times 20 \times e^{-4}}{1.4e^{-4}(1 - 4)} = 22.8 \text{ hrs.}$$

Note: Newton's method is based on the equation of the tangent: $y - y_1 = f'(x_1)(x - x_1)$.

Usually we find the approximation when $y = 0$.

Here, $y = 0.3$

$$\therefore x - x_1 = \frac{y - y_1}{f'(x_1)}$$

$$x = x_1 + \frac{y - y_1}{f'(x_1)}.$$

Question 14

(a) (i) $CS \perp AD$ (semi-circle angle)

For the same reason, $CT \perp BT$ and $AD \perp DB$

$\therefore CTDS$ is a rectangle (three right angles)

(ii) $MS = MC$ (= radii)

$XS = XC$ (in a rectangle, the diagonals bisect)

MX is common

$\therefore \triangle MXS \cong \triangle MXC$ (SSS)

(iii) $\angle MSX = \angle MCX$ (corresponding angles in congruent triangles)

$\therefore \angle MSX = 90^\circ$

$\therefore ST$ is a tangent (tangent is perpendicular to the radius at the point of contact).

(b) (i) At maximum height, $\dot{y} = 0$

$$70 \sin \theta - 9.8t = 0$$

$$t = \frac{70 \sin \theta}{9.8} \text{ s.}$$

$$y = 70t \sin \theta - 4.9t^2$$

$$= \frac{4900 \sin^2 \theta}{9.8} - \frac{4.9 \times 4900 \sin^2 \theta}{9.8^2}$$

$$= 250 \sin^2 \theta.$$

(ii) Substitute $t = \frac{70 \sin \theta}{9.8}$ into $x = 70t \cos \theta$

$$x = \frac{4900 \sin \theta \cos \theta}{9.8} = 500 \sin \theta \cos \theta$$

$$= 250 \sin 2\theta.$$

(iii) Solving $250 \sin^2 \theta \geq 150$ gives $\sin^2 \theta \geq \frac{3}{5}$


$$\sin \theta \geq \frac{\sqrt{15}}{5}, \therefore 50.8^\circ \leq \theta \leq 129.2^\circ. \quad (1)$$

Solving $125 \leq 250 \sin 2\theta \leq 180$ gives

$$\frac{1}{2} \leq \sin 2\theta \leq \frac{18}{25}$$

$$30^\circ + k.360^\circ \leq 2\theta \leq 46^\circ + k.360^\circ$$

$$\text{or } 134^\circ + k.360^\circ \leq 2\theta \leq 150^\circ + k.360^\circ.$$



Dividing by 2,

$$15^\circ + k180^\circ \leq \theta \leq 23^\circ + k180^\circ$$

$$\text{or } 67^\circ + k180^\circ \leq \theta \leq 75^\circ + k180^\circ.$$

\therefore To satisfy (1), $67^\circ \leq \theta \leq 75^\circ$.

(c) (i) $\cos \alpha = \frac{BG}{u}, \therefore BG = u \cos \alpha.$

$$\sin \alpha = \frac{PG}{u}, \therefore PG = u \sin \alpha.$$

$$AG^2 = BG^2 + 1^2 - 2BG \cos 60^\circ$$

$$= BG^2 + 1 - BG = u^2 \cos^2 \alpha + 1 - u \cos \alpha.$$

$$r^2 = PG^2 + AG^2$$

$$= u^2 \sin^2 \alpha + u^2 \cos^2 \alpha + 1 - u \cos \alpha$$

$$= u^2 (\sin^2 \alpha + \cos^2 \alpha) + 1 - u \cos \alpha$$

$$= u^2 + 1 - u \cos \alpha.$$

$$\therefore r = \sqrt{1 + u^2 - u \cos \alpha}.$$

(ii) $\frac{dr}{dt} = \frac{dr}{du} \frac{du}{dt} = \frac{2u - \cos \alpha}{2\sqrt{1 + u^2 - u \cos \alpha}} \frac{du}{dt}$

When $t = 5$ min., $u = 30$ km, $\frac{du}{dt} = 6$ km/min.,

$$\frac{dr}{dt} = \frac{60 - \cos \alpha}{2\sqrt{901 - 30 \cos \alpha}} \times 6$$

$$= \frac{180 - 3 \cos \alpha}{\sqrt{901 - 30 \cos \alpha}} \text{ km/min.}$$